sam.nitk.ac.in sam@nitk.edu.in

Computational Mathematics - MA 608 **Problem Sheet - 2**

Curve Fitting

1. Convert the following equations into linear form

(a)
$$y = \frac{x}{a+bx}$$

(c)
$$xa^y = b$$

(e)
$$xy = ax + b$$

(g)
$$y = ax + bxy$$
.

(b)
$$y = \frac{ax+b}{x}$$

(c)
$$xa^{y} = b$$
 (e) $xy = ax + b$
(d) $y = \frac{b}{x(x-a)}$ (f) $y = ax + bx^{2}$

$$(f) y = ax + bx^2$$

2. Find the most plausible values of *x*, *y* and *z* from the equations

$$x + 3y - 3z = -14$$

$$4x + y + 4z = 21$$

$$3x + 2y - 5z = 5$$

$$x - y + 2z = 3$$

by forming the normal equations.

- 3. Construct a least squares quadratic approximation to the function $y(x) = \sin x$ on $[0, \pi/2]$ with respect to the weight function W(x) = 1.
- 4. If *P* is the pull required to lift a load *W* by means of a pulley block, find a linear law of the form P = mW + C, connecting P and W, using the following data.

P (in kg.)	12	15	21	25
W (in kg.)	50	70	100	120

Compute *P* when W = 150 kg.

5. By the method of least squares, find the straight line that best fits the following data.

x	1	2	3	4	5
Ŋ	14	27	40	55	68

6. Fit a straight line to the following data and estimate the value of y corresponding to x = 6.

$\boldsymbol{\mathcal{X}}$	0	5	10	15	20	25
y	12	15	17	22	24	30

7. Fit a second degree parabola by taking x as the independent variable.

х	1	1	2	3	4
y	1	5	10	22	38

8. Using the method of least squares, fit a curve of the form $y = ab^x$ to the following data.

χ	:	1	2	3	4
1	,	4	11	35	100

9. Fit a curve of the form $y = ab^x$ to the following data.

Year (x)	1951	1952	1953	1954	1955	1956	1957
Production							
in tone (y)	201	263	314	395	427	504	612

10. Fit a curve of the form $y = ax^b$ for the following data, where a and b are constants.

х	61	26	7	10
у	350	400	500	600

11. Using the principle of least squares, fit an equation of the form $y = ae^{bx}$ (a > 0) to the data.

	х	1	2	3	4
ĺ	у	1.65	2.7	4.5	7.35

12. The pressure and volume of a gas are related by the equation $pc^{\lambda} = k$ (λ and k are constants). Fit this equation for the following data, using the principle of least squares.

p	0.5	1.0	1.5	2.0	2.5	3.0
v	1.62	1.00	0.75	0.62	0.52	0.46

13. Two quantities of x and y are measured and corresponding values are given in the following table.

				60			
Ī	y	5.5	9.1	14.9	22.8	33.3	46

Find a second degree parabola to the data.
